

Fermionic fields in the d -dimensional anti-de Sitter spacetime

R.R. Metsaev

*Department of Theoretical Physics, P.N. Lebedev Physical Institute, Leninsky prospect
53, 117924, Moscow, Russia*

Abstract

Arbitrary spin free massless fermionic fields corresponding to mixed symmetry representations of the $SO(d-1)$ compact group and propagating in even d -dimensional anti-de Sitter spacetime are investigated. Free wave equations of motion, subsidiary conditions and the corresponding gauge transformations for such fields are proposed. The lowest eigenvalues of the energy operator for the massless fields and the gauge parameter fields are derived. The results are formulated in $SO(d-1, 2)$ covariant form as well as in terms of intrinsic coordinates.

hep-th/9802097, published in *Phys. Lett. B* 419 (1998) 49

In view of the aesthetic features of anti-de Sitter field theory a interest in this theory was periodically renewed (see [1]-[8] and references therein). One of the interesting directions of this theory is the higher-spin massless field theory. At present there is reason for revival of interest in higher-spin massless fields. Recently it was discovered [9] that the consistent equations of motion of interacting higher-spin massless fields in four dimensional ($d = 4$) anti-de Sitter spacetime could be set up with the help of higher-spin superalgebras (for review see [7]). These equations for the case of higher spacetime dimensions $d > 4$ have been generalized in [10]. Because these equations are formulated in terms of wavefunctions which depend on usual spacetime coordinates and certain twistor variables it is not clear immediately what kind of fields they describe. In [9], it was established that in four dimensions ($d = 4$) they describe unitary dynamics of massless fields of all spins in anti-de Sitter spacetime.

The long-term motivation of our investigation is to provide an answer to question: do the higher dimensional theories of Ref.[10] describe higher-spin massless fields? The first step in such an investigation is a description of free equations of motion for arbitrary spin massless fields. In contrast to completeness of description for $d = 4$ ([1],[2],[8],[11]-[13]), not much is known on the higher-spin fermionic massless fields for $d > 4$ even at the level of free fields, unless considerations are restricted to totally (anti)symmetric representations ([14]-[16]). The present paper is a sequel to our paper [17] where description of bosonic massless fields of all spins in anti-de Sitter spacetime of arbitrary dimensions was developed. The case of half-integral spins, presented here is a necessary step in our study of massless fields of all spins for arbitrary d . With higher-spin theories in mind we also hope that our results may be of wider interest, especially to anti-de Sitter supergravity theories.

Let us first formulate the main problems we solve in this letter. A positive-energy lowest weight irreducible representation of the $SO(d-1, 2)$ group denoted as $D(E_0, \mathbf{m})$, is defined by E_0 , the lowest eigenvalue of the energy operator, and by $\mathbf{m} = (m_1, \dots, m_\nu)$, $\nu \equiv \frac{d-2}{2}$, which is the weight of the unitary representation of the $SO(d-1)$ group. The m_i are half integers for fermions. For the case of $SO(3, 2)$ group it has been discovered [11] that fermionic massless fields propagating in four dimensional anti-de Sitter spacetime are associated with $D(s+1, s)$ i.e. spin- s anti-de Sitter massless particle takes lowest value of energy equal to $E_0 = s+1$. In [16] for the case of arbitrary d it was found that $E_0 = s+d-3$ for totally symmetric massless representation $\mathbf{m}_{sym} = (s, 1/2, \dots, 1/2)$, $s \geq 3/2$, and $E_0 = d-s'-1/2$ for the totally antisymmetric one $\mathbf{m}_{as} = (3/2, \dots, 3/2, 1/2 \dots 1/2)$ (where the $3/2$ occurs $s' \leq \nu$ times in this sequence). The cases $\mathbf{m}_{sym}$ and $\mathbf{m}_{as}$ are very special: there are many representations corresponding to arbitrary $\mathbf{m}$ (mixed symmetry representations). In this paper we construct free equations of motion and gauge transformations for massless fields with arbitrary $\mathbf{m}$ and determine the E_0 . Here we restrict ourselves to the case of even spacetime dimension d .

Let us describe our conventions and notation. We describe the anti-de Sitter spacetime as a hyperboloid

$$\eta_{AB} y^A y^B = -1, \quad \eta_{AB} = (-, +, \dots, +, -), \quad A, B = 0, 1, \dots, d-1, d+1, \quad (1)$$

in $d+1$ - dimensional pseudo-Euclidean space with metric tensor η_{AB} . The indices A, B are raised and lowered by η^{AB} and η_{AB} respectively. In what follows to simplify our

expressions we will drop the metric tensor η_{AB} in scalar products. The generators J^{AB} of the $SO(d-1, 2)$ group satisfy the commutation relations

$$[J^{AB}, J^{CD}] = \eta^{BC} J^{AD} + 3 \text{ terms}.$$

As is usual, we split J^{AB} into an orbital part L^{AB} and a spin part M^{AB} : $J^{AB} = L^{AB} + M^{AB}$. The realization of L^{AB} in terms of differential operators defined on the hyperboloid (1) is:

$$L^{AB} = y^A \nabla^B - y^B \nabla^A, \quad \nabla^A \equiv \theta^{AB} \frac{\partial}{\partial y^B}, \quad \theta^{AB} \equiv \eta^{AB} + y^A y^B,$$

The tangent derivative ∇^A has the properties $y^A \nabla^A = 0$, $\nabla^A y^A = d$ and satisfies the commutation relations

$$[\nabla^A, y^B] = \theta^{AB}, \quad [\nabla^A, \nabla^B] = -L^{AB}.$$

A form for M^{AB} depends on the realization of the representations. We will use the tensor realization of representations. As the carriers for $D(E_0, \mathbf{m})$ we use tensor-spinor field

$$\Psi^{A(\mathbf{m})} = \Psi^{A_1^1, \dots, A_1^{m'_1}, \dots, A_\nu^1, \dots, A_\nu^{m'_\nu}}, \quad m'_i \equiv m_i - \frac{1}{2},$$

defined on the hyperboloid (1). The Ψ has one spinor index which we do not show explicitly. By definition, $\Psi^{A(\mathbf{m})}$ is a tensor-spinor field whose $d+1$ spacetime indices A have the structure of the Young tableaux (YT) corresponding to the irreps of the $SO(d-1, 2)$ group labeled by $\mathbf{m}' = (m'_1 \dots m'_\nu)$. In what follows we use the notation $\text{YT}(\mathbf{m}')$ to indicate such YT. At the same time the $\mathbf{m}$ is the weight of a representation of the $SO(d-1)$ group, and the m_i satisfy the inequalities

$$m_1 \geq \dots \geq m_\nu \geq 1/2. \quad (2)$$

In the language of YT, the m'_i indicates the number of boxes in the i -th row of $\text{YT}(\mathbf{m}')$. Thus given an arbitrary lowest weight massless representation $D(E_0, \mathbf{m})$ we assume that a covariant description can be formulated with the field $\Psi^{A(\mathbf{m})}$. To avoid cumbersome tensor expressions we introduce ν creation and annihilation operators a_l^A and $\bar{a}_l^A$ ($l = 1, \dots, \nu$, $\nu = (d-2)/2$) which satisfy

$$[\bar{a}_i^A, a_j^B] = \eta^{AB} \delta_{ij}, \quad \bar{a}_i^A |0\rangle = 0$$

and construct a Fock space vector

$$|\Psi\rangle \equiv \prod_{l=1}^{\nu} \prod_{i_l=1}^{m'_l} a_l^{A_l^{i_l}} \Psi^{A_1^1, \dots, A_1^{m'_1}, \dots, A_\nu^1, \dots, A_\nu^{m'_\nu}} |0\rangle. \quad (3)$$

For a realization of this kind, M^{AB} has the form

$$M^{AB} = M'^{AB} + \sigma^{AB},$$

where

$$M'^{AB} = \sum_{l=1}^{\nu} (a_l^A \bar{a}_l^B - a_l^B \bar{a}_l^A), \quad \sigma^{AB} = \frac{1}{4} (\gamma^A \gamma^B - \gamma^B \gamma^A)$$

and γ^A are the gamma matrices: $\{\gamma^A, \gamma^B\} = 2\eta^{AB}$. Throughout of the paper, unless otherwise specified, the indices i, j, l, n run over $1, \dots, \nu$. For these indices we drop the summation over repeated indices.

If the $\Psi^{A(\mathbf{m})}$ is associated with $\mathbf{YT}(\mathbf{m}')$ then the $|\Psi\rangle$ satisfies

$$(a_{ii} - m'_i)|\Psi\rangle = 0, \quad a_{ij}^-|\Psi\rangle = 0, \quad \varepsilon^{ij}a_{ij}|\Psi\rangle = 0, \quad (4)$$

where in (4) and below we use the notation

$$a_{ij} \equiv a_i^A \bar{a}_j^A, \quad a_{ij}^- \equiv \bar{a}_i^A \bar{a}_j^A, \quad a_{ij}^+ \equiv a_i^A a_j^A,$$

and $\varepsilon^{ij} = 1(0)$ for $i < j(i \geq j)$. The 1st equation in (4) tells us that a_i occurs m'_i times on the right hand side of eq.(3). Tracelessness of $\Psi^{A(\mathbf{m})}$ is reflected in the 2nd equation in (4). The 3rd equation in (4) is a consequence of the following Young's symmetrization rule we use: (i) first we perform alternating with respect to indices in all columns, (ii) then we perform symmetrization with respect to indices in all rows (for more explanations see [19])

Being the carrier for the $D(E_0, \mathbf{m})$, the field $|\Psi\rangle$ satisfies the equation

$$(Q - \langle Q \rangle)|\Psi\rangle = 0, \quad Q \equiv \frac{1}{2} J^{AB} J^{AB} \quad (5)$$

and allows the following subsidiary covariant constraints

$$\bar{\nabla}_n |\Psi\rangle = 0, \quad (\text{divergencelessness}) \quad (6)$$

$$\bar{y}_n |\Psi\rangle = 0, \quad (\text{transversality}) \quad (7)$$

$$\bar{\gamma}_n |\Psi\rangle = 0, \quad (8)$$

$n = 1, \dots, \nu$. In (5) Q is the second order Casimir operator of the $so(d-1, 2)$ algebra while $\langle Q \rangle$ is its eigenvalue for $D(E_0, \mathbf{m})$

$$\langle Q \rangle = -E_0(E_0 + 1 - d) - \sum_{l=1}^{\nu} m_l(m_l - 2l + d - 1),$$

where E_0 is the lowest eigenvalue of $\mathbf{i}J^{d+10}$. The $\langle Q \rangle$ can be calculated according to the well known procedure [18]. In (6)-(8) and below we use the notation

$$\nabla_n \equiv \mathbf{a}_n^A \nabla^A, \quad \bar{\nabla}_n \equiv \nabla^A \bar{\mathbf{a}}_n^A, \quad \mathbf{a}_n^A \equiv \theta^{AB} a_n^B, \quad \bar{\mathbf{a}}_n^A \equiv \theta^{AB} \bar{a}_n^B.$$

$$y_n \equiv y^A a_n^A, \quad \bar{y}_n \equiv y^A \bar{a}_n^A, \quad \gamma_n \equiv \gamma^A a_n^A, \quad \bar{\gamma}_n \equiv \gamma^A \bar{a}_n^A.$$

Introducing

$$\mathbf{a}_{ij} \equiv \mathbf{a}_i^A \bar{\mathbf{a}}_j^A, \quad \mathbf{a}_{ij}^- \equiv \bar{\mathbf{a}}_i^A \bar{\mathbf{a}}_j^A, \quad \mathbf{a}_{ij}^+ \equiv \mathbf{a}_i^A \mathbf{a}_j^A$$

and using the constraints (7) and the equalities

$$a_{ij} = \mathbf{a}_{ij} - y_i \bar{y}_j, \quad a_{ij}^- = \mathbf{a}_{ij}^- - \bar{y}_i \bar{y}_j, \quad (9)$$

the constraints (4) can be rewritten in the form which we use in what follows:

$$(\mathbf{a}_{ii} - m'_i)|\Psi\rangle = 0, \quad \mathbf{a}_{ij}^-|\Psi\rangle = 0, \quad \varepsilon^{ij}\mathbf{a}_{ij}|\Psi\rangle = 0. \quad (10)$$

For E_0 corresponding to the massless case a solution of Eqs. (5)-(8) decomposes into a physical massless representations and some additional representations corresponding to pure gauge fields (i.e. the situation is similar to the $d = 4$ case [12]).

Now rewriting the expression for Q as

$$Q = -\nabla^2 + M^{AB}L^{AB} + \frac{1}{2}M^{AB}M^{AB}, \quad \nabla^2 \equiv \nabla^A \nabla_A$$

and taking into account the easily derived equalities (using eqs.(6)-(8) and (10))

$$\sigma^{AB}L^{AB}|\Psi\rangle = \not{y} \not{\nabla} |\Psi\rangle, \quad M'^{AB}M'^{AB}|\Psi\rangle = -2 \sum_{l=1}^{\nu} m'_l(m'_l - 2l + d + 1)|\Psi\rangle,$$

$$M'^{AB}L^{AB}|\Psi\rangle = 2 \sum_{l=1}^{\nu} m'_l |\Psi\rangle, \quad M'^{AB}\sigma^{AB}|\Psi\rangle = - \sum_{l=1}^{\nu} m'_l |\Psi\rangle,$$

$\not{\nabla}^2 = \nabla^2 - \not{y} \not{\nabla}$, where $\not{\nabla} \equiv \gamma^A \nabla_A$, $\not{y} \equiv \gamma^A y_A$, $\{\not{\nabla}, \not{y}\} = d$, we can rewrite Eq. (5) as

$$(\not{\nabla}^2 - (E_0 + \frac{1}{2})(E_0 + \frac{1}{2} - d))|\Psi\rangle = 0. \quad (11)$$

To define E_0 corresponding to massless fields we construct gauge invariant equations of motion for $|\Psi\rangle$. In order to formulate gauge transformations we use the gauge parameters fields whose spacetime indices correspond to the YT which one can make by removing one box from the $YT(\mathbf{m}')$. Thus the most general gauge transformations we start with are

$$\delta_{(n)}|\Psi\rangle \sim \nabla_n|\Lambda_n\rangle + y_n|R_n\rangle + \gamma_n|S_n\rangle, \quad (12)$$

where the gauge parameters fields $|\Lambda_n\rangle$, $|R_n\rangle$ and $|S_n\rangle$ are associated with $YT(\mathbf{m}'_{(n)})$, and i -th component of the $\mathbf{m}'_{(n)}$ is equal to $m'_{i(n)} = m'_i - \delta_{in}$. The $YT(\mathbf{m}'_{(n)})$ is obtained by removing one box from n -th row of the $YT(\mathbf{m}')$. We assume that only those $|\Lambda_n\rangle$, $|R_n\rangle$ and $|S_n\rangle$ are non-zero whose $\mathbf{m}'_{(n)}$ satisfy the inequalities

$$m'_{1(n)} \geq \dots \geq m'_{\nu(n)} \geq 0. \quad (13)$$

Given the $\mathbf{m}'$, the set of those n whose $\mathbf{m}'_{(n)}$ satisfy (13) will be referred to as $\mathbf{S}(\mathbf{m}')$, while the number of such n will be referred to as ν' . Thus we have ν' nontrivial gauge transformations. We impose on the $|\Lambda_n\rangle$, $|R_n\rangle$ and $|S_n\rangle$ the constraints similar to (6)-(8)

$$\bar{\nabla}_i|\Lambda_n\rangle = 0, \quad \bar{y}_i|\Lambda_n\rangle = 0, \quad \bar{\gamma}_i|\Lambda_n\rangle = 0, \quad (14)$$

and constraints obtained from (14) by replacing $\Lambda \rightarrow R$ and $\Lambda \rightarrow S$. Since $|\Lambda_n\rangle$, $|R_n\rangle$ and $|S_n\rangle$ correspond to $\mathbf{YT}(\mathbf{m}'_{(n)})$, they satisfy the constraints

$$(a_{ii} - m'_{i(n)})|\Lambda_n\rangle = 0, \quad a_{ij}^-|\Lambda_n\rangle = 0, \quad \varepsilon^{ij}a_{ij}|\Lambda_n\rangle = 0, \quad (15)$$

and those which are obtainable from (15) by replacing $\Lambda \rightarrow R$ and $\Lambda \rightarrow S$. Using (9) and (14), the constraints (15) can be rewritten in the form similar to (10)

$$(\mathbf{a}_{ii} - m'_{i(n)})|\Lambda_n\rangle = 0, \quad \mathbf{a}_{ij}^-|\Lambda_n\rangle = 0, \quad \varepsilon^{ij}\mathbf{a}_{ij}|\Lambda_n\rangle = 0. \quad (16)$$

Now we should find such $\delta_{(n)}|\Psi\rangle$ and E_0 that the constraints and the equations of motion (6)-(8), (10), (11) be invariant with respect to gauge transformations. We proceed in the following way.

(i) from the invariance requirement of (7) (i.e. $\bar{y}_i\delta_{(n)}|\Psi\rangle = 0$) and (12), (14) one gets

$$|R_n\rangle = -\sum_{l=1}^{\nu} \mathbf{a}_{ln}|\Lambda_l\rangle + \not{y}|S_n\rangle. \quad (17)$$

By substituting (17) into (12) we obtain

$$\delta_{(n)}|\Psi\rangle \sim D_n|\Lambda_n\rangle + (\gamma\mathbf{a}_n)|S_n\rangle, \quad n \in \mathbf{S}(\mathbf{m}'), \quad (18)$$

where $(\gamma\mathbf{a}_n) \equiv \gamma^A\mathbf{a}_n^A$ and

$$D_n \equiv \nabla_n + \sum_{l=1}^{\nu} (-y_l\mathbf{a}_{nl} + \mathbf{a}_{nl}^+\bar{y}_l). \quad (19)$$

In (19) we add by hand the 2nd term in the sum. Due to (14) the gauge transformations (18) are unaffected by this term. The D_n introduced in such a way allows the representation $D_n = a_n^A y^B J'^{AB}$, where $J'^{AB} \equiv L^{AB} + M'^{AB}$, which is very convenient in practical calculations. Some useful commutation relations are

$$\begin{aligned} [\mathbf{a}_{ij}, D_n] &= \delta_{jn}D_i, & [\mathbf{a}_{ij}^-, D_n] &= \delta_{in}\bar{D}_j + \delta_{jn}\bar{D}_i, \\ [\mathbf{a}_{ij}, \mathbf{a}_{nl}] &= \delta_{jn}\mathbf{a}_{il} - \delta_{il}\mathbf{a}_{nj}, & [\mathbf{a}_{ij}^-, \mathbf{a}_{nl}] &= \delta_{jn}\mathbf{a}_{il}^- + \delta_{in}\mathbf{a}_{jl}^-, \end{aligned}$$

where $\bar{D}_n \equiv J'^{AB}\bar{a}_n^A y^B$. The usage of D_i , $\bar{D}_i$, $\mathbf{a}_{ij}^\pm$ and $\mathbf{a}_{ij}$ is advantageous since all of them commute with y_n and $\bar{y}_n$.

(ii) using (14), (16) one can verify that the 1st and 2nd constraints from (10) are invariant with respect to (18) i.e. the equations $(\mathbf{a}_{ii} - m'_i)\delta_{(n)}|\Psi\rangle = 0$ and $\mathbf{a}_{ij}^-\delta_{(n)}|\Psi\rangle = 0$ are fulfilled. The invariance requirement of 3rd constraint from (10) (i.e. $\varepsilon^{ij}\mathbf{a}_{ij}\delta_{(n)}|\Psi\rangle = 0$) is the most difficult point. Fortunately, this part of analysis is similar to that of the bosonic case ([17]). Our result for $\delta_{(n)}$ is:

$$\begin{aligned} \delta_{(n)}|\Psi\rangle &= \mathcal{D}_n|\Lambda_n\rangle + \mathcal{G}_n|S_n\rangle, \\ \mathcal{D}_n &\equiv \sum_{l=1}^{\nu} P_{nl}D_l, & \mathcal{G}_n &\equiv \sum_{l=1}^{\nu} P_{nl}(\gamma\mathbf{a}_l), \end{aligned} \quad (20)$$

where

$$P_{nl} = \sum_{j=0}^{n-1} (-)^j \sum_{l_1 \dots l_{j+1}=1}^{\nu} \delta_{nl_{j+1}} \delta_{l_1 l} \prod_{i=1}^j \frac{\varepsilon^{l_i l_{i+1}}}{\lambda_{l_i n}} \mathbf{a}_{l_{i+1} l_i}, \quad (21)$$

$$\lambda_{ln} \equiv m_l - m_n + n - l + 1.$$

In (21) the $\mathbf{a}_{l_{i+1} l_i}$ are ordered as follows: $\mathbf{a}_{l_{j+1} l_j} \dots \mathbf{a}_{l_2 l_1}$. As an illustration of (20) we write down $\delta_{(n)}|\Psi\rangle$ for $n = 1, 2, 3$ assuming that such an n belongs to $\mathbf{S}(\mathbf{m}')$:

$$\begin{aligned} \delta_{(1)}|\Psi\rangle &= D_1|\Lambda_1\rangle + (\gamma\mathbf{a}_1)|S_1\rangle, \\ \delta_{(2)}|\Psi\rangle &= D_2|\Lambda_2\rangle + (\gamma\mathbf{a}_2)|S_2\rangle - \frac{\mathbf{a}_{21}}{\lambda_{12}}(D_1|\Lambda_2\rangle + (\gamma\mathbf{a}_1)|S_2\rangle), \\ \delta_{(3)}|\Psi\rangle &= D_3|\Lambda_3\rangle + (\gamma\mathbf{a}_3)|S_3\rangle - \frac{\mathbf{a}_{31}}{\lambda_{13}}(D_1|\Lambda_3\rangle + (\gamma\mathbf{a}_1)|S_3\rangle) - \frac{\mathbf{a}_{32}}{\lambda_{23}}(D_2|\Lambda_3\rangle + (\gamma\mathbf{a}_2)|S_3\rangle) \\ &\quad + \frac{\mathbf{a}_{32}\mathbf{a}_{21}}{\lambda_{13}\lambda_{23}}(D_1|\Lambda_3\rangle + (\gamma\mathbf{a}_1)|S_3\rangle). \end{aligned}$$

(iii) from the invariance requirement of eq.(8) (i.e. $\bar{\gamma}_i \delta_{(n)}|\Psi\rangle = 0$) we get the equation for $|\Lambda_n\rangle$ and $|S_n\rangle$

$$(\nabla - (m'_n - n)\psi)|\Lambda_n\rangle + (d + 2(m'_n - n))|S_n\rangle = 0 \quad (22)$$

and from the invariance requirement of eq.(6) (i.e. $\bar{\nabla}_i \delta_{(n)}|\Psi\rangle = 0$) we get the equation

$$(\nabla^2 - (m'_n - n)(m'_n - n - 1 + d))|\Lambda_n\rangle + (\nabla + (m'_n - n + d)\psi)|S_n\rangle = 0. \quad (23)$$

Finally from the invariance requirement of eq.(11) (i.e. $(\nabla^2 - \dots)\delta_{(n)}|\Psi\rangle = 0$) and from Eqs.(22), (23) we derive the equation for E_0

$$(E_0 + \frac{1}{2})(E_0 + \frac{1}{2} - d) = (m'_n - n - 1)(m'_n - n - 1 + d), \quad n \in \mathbf{S}(\mathbf{m}'), \quad (24)$$

whose solutions read as

$$E_{0(n)}^{(1)} = m_n - n - 2 + d, \quad E_{0(n)}^{(2)} = n + 1 - m_n, \quad n \in \mathbf{S}(\mathbf{m}'). \quad (25)$$

As seen from (25) there exists an arbitrariness of E_0 parametrized by subscript n which labels gauge transformations and by superscripts (1), (2) which label two solutions of quadratic equation (24). Because the values of E_0 have been derived by exploiting gauge invariance we can conclude that the gauge invariance by itself does not uniquely determine the physical relevant value of E_0 (in this regard the situation is similar to the $d = 4$ case, see [1]). To choose physical relevant value of E_0 we exploit the unitarity condition, that is: 1) hermiticity $(iJ^{AB})^\dagger = iJ^{AB}$; 2) the positive norm requirement. The resulting procedure is the same that has been previously encountered in the bosonic case [17]. Therefore we do not give deal of technical details and formulate the result.

Given $\text{YT}(\mathbf{m}')$ let $k, k = 1 \dots \nu$, indicates maximal number of upper rows which have the same number of boxes. We call such Young tableaux the level- k $\text{YT}(\mathbf{m}')$. For the case of level- k Young tableaux the inequalities (2) can be rewritten as

$$m_1 = \dots = m_k > m_{k+1} \geq m_{k+2} \geq \dots \geq m_\nu \geq 1/2. \quad (26)$$

Then making use of unitarity condition one proves [17] that for the level- k Young tableaux the E_0 should satisfy the inequality

$$E_0 \geq m_k - k - 2 + d. \quad (27)$$

Comparing (25) with (27) we conclude that only $E_{0(n=k)}^{(1)}$ satisfies the unitarity condition. Thus anti-de Sitter fermionic massless particles described by level- k $\text{YT}(\mathbf{m}')$ takes lowest value of energy equal to

$$E_0 = m_k - k - 2 + d. \quad (28)$$

Note that it is gauge transformation with $n = k$ (20) that leads to relevant E_0 , i.e. given level- k $\text{YT}(\mathbf{m}')$ only the gauge transformation $\delta_{(k)}$ respects the unitarity. Therefore only the $\delta_{(k)}$ will be used in what follows. From now on we use letter k to indicate level of $\text{YT}(\mathbf{m}')$. Our result for E_0 (28) cannot be extended to cover the case $\mathbf{m}' = 0$ because the E_0 obtained are relevant only for gauge fields. This case should be considered in its own right and the relevant value $E_0(\text{for } \mathbf{m}' = 0) = (d - 1)/2$ can be obtained by using requirement of conformal invariance (see [21]). The case $\mathbf{m}' = 0, d = 4$ has been investigated in [22].

Up to now we analysed second-order equations of motion for Ψ and gauge invariance of these equations. Now we would like to derive first-order equations of motion and corresponding gauge transformations. To do that we use the relations

$$\nabla^2 = \kappa^2 - d\kappa, \quad \kappa \equiv \sigma^{AB} L^{AB}, \quad \kappa = \not{y} \not{\nabla}, \quad (29)$$

and rewrite equation of motions (11) as follows:

$$(\kappa - E_0 - \frac{1}{2})(\kappa + E_0 + \frac{1}{2} - d)|\Psi\rangle = 0.$$

Thus we could use the following first-order equations of motion:

$$(\kappa + E_0 + \frac{1}{2} - d)|\Psi\rangle = 0, \quad (30)$$

$$(\kappa - E_0 - \frac{1}{2})|\tilde{\Psi}\rangle = 0. \quad (31)$$

Due to relation

$$(\kappa + E_0 + \frac{1}{2} - d)|\Psi\rangle = \not{y}(\kappa - E_0 - \frac{1}{2})\not{y}|\Psi\rangle$$

it is clear that $|\Psi\rangle$ and $|\tilde{\Psi}\rangle$ are related by $|\Psi\rangle = \not{y}|\tilde{\Psi}\rangle$, i.e. equations (30) and (31) are equivalent. We will use the equation of motion given by (30). Now we should verify

gauge invariance of the first-order equation (30) with respect to gauge transformations (20). It turns out that the invariance requirement of (30) with respect to (20) leads to

$$|S_k\rangle = 0. \quad (32)$$

Thus because of (26) and (32) the final form of gauge transformations is

$$\delta_{(k)}|\Psi\rangle = \sum_{j=0}^{k-1} (-)^j \sum_{l_1 \dots l_{j+1}=1}^{\nu} \delta_{kl_{j+1}} \prod_{i=1}^j \frac{\varepsilon^{l_i l_{i+1}} \mathbf{a}_{l_{i+1} l_i}}{k+1-l_i} D_{l_1} |\Lambda_k\rangle. \quad (33)$$

The equations of motion for Λ can be obtained from (22) and (32)

$$(\nabla - (m_k - \frac{1}{2} - k) \not{y}) |\Lambda_k\rangle = 0 \quad (34)$$

which can also be rewritten in terms of κ and E_0^Λ

$$(\kappa + E_0^\Lambda + \frac{1}{2} - d) |\Lambda_k\rangle = 0, \quad (35)$$

where we introduce lowest energy value for $|\Lambda_k\rangle$: $E_0^\Lambda = E_0 + 1$

$$E_0^\Lambda = m_k - k - 1 + d. \quad (36)$$

With the values for E_0^Λ at hand we are ready to provide an answer to the question: do the gauge parameter fields meet the masslessness criteria? Because the inter-relation between of spin $\mathbf{m}$ and energy value E_0 for massless field is given by (28) we should express the E_0^Λ in terms of $\mathbf{m}^\Lambda$ and k^Λ , where k^Λ is a level of $\mathbf{YT}(\mathbf{m}^\Lambda)$. Due to relations $k^\Lambda = k - 1$, $m_{k^\Lambda}^\Lambda = m_k - \delta_{k1}$ we transform (36) to

$$E_0^\Lambda = m_{k^\Lambda}^\Lambda - k^\Lambda - 2 + d + \delta_{k1}.$$

Comparing this relation with (28) we conclude that only for $k > 1$ the gauge parameters are massless fields while for $k = 1$ they are massive fields.

Thus we have constructed equations of motion (30) which respect gauge transformations (33), where the gauge parameter fields Λ satisfy the constraints (14), (16) and equations of motion (35). The relevant values of E_0 and E_0^Λ are given by (28) and (36). Note that the equations of motion are written in terms of the operator κ (see equation (29)) introduced in [20] while constructing the equation of motion for the field associated with the representation $D(E_0, 1/2)$. The κ is expressible in terms of the orbital momentum L^{AB} (see equation (29)). Now we would like to rewrite our equations of motion in terms of complete angular momentum J^{AB} . We believe that such a formulation will form a good basis for establishing an action leading to the equations of motion under consideration. To this end let us first multiply equations (30) and (35) by $\not{y}$. Then making use of equalities

$$\gamma^A y^B J^{AB} = \not{y} + \sum_{l=1}^{\nu} (\gamma_l \bar{y}_l - y_l \bar{\gamma}_l) + \frac{1}{2} d \not{y}, \quad (37)$$

$$a_n^A y^B J^{AB} = D_n + \frac{1}{2} a_n^A \theta^{AB} \gamma^B \not{y}, \quad (38)$$

and the constraints (7), (8), (14) we get the desired form for equations of motion:

$$\left(\gamma^A y^B J^{AB} - \left(E_0 - \frac{d-1}{2}\right) \not{y}\right) |\Psi\rangle = 0, \quad (39)$$

$$\left(\gamma^A y^B J^{AB} - \left(E_0^\Lambda - \frac{d-1}{2}\right) \not{y}\right) |\Lambda_k\rangle = 0, \quad (40)$$

where E_0 and E_0^Λ are given by (28) and (36). Because the term $D_{l_1} |\Lambda_k\rangle$ from gauge transformations (33) can be expressed as

$$D_l |\Lambda_k\rangle = \left(a_l^A - \frac{a_l^C \theta^{CD} \gamma^D}{2E_0 + 3 - d} \gamma^A\right) y^B J^{AB} |\Lambda_k\rangle, \quad (41)$$

it is seen that the gauge transformations (33) are also expressible in terms of J^{AB} . Note that in deriving of (40) and (41) it is necessary to use the equations of motion for $|\Lambda_k\rangle$ (35) and the relevant values of E_0 and E_0^Λ given by (28) and (36).

Now we would like to transform our results to intrinsic coordinates, in terms of covariant derivatives and vierbein fields. Let x^μ , $\mu = 0, 1, \dots, d-1$ be the intrinsic coordinates for anti-de Sitter spacetime and let $y^A(x)$ be imbedding map, where $y^A(x)$ satisfy (1). The relationship between $d+1$ -dimensional tensor-spinor field $\Psi^{A\dots}$ and the usual Rarita-Schwinger tensor-spinor field $\psi_{\mu\dots}$ is given by (see [12])

$$\psi_\mu \dots(x) = M^{-1}(x) y_\mu^A \dots \Psi^{A\dots}(y),$$

where $y_\mu^A \equiv \partial_\mu y^A$ and M is an $2^{d/2} \times 2^{d/2}$ matrix defined by

$$M^{-1} \partial_\mu M = \frac{1}{2} \omega_\mu^{ab} \sigma_{ab} + \frac{1}{2} e_\mu^a \gamma_a, \quad M^{-1} (\not{y} \partial_\mu \not{y}) M = e_\mu^a \gamma_a, \quad (42)$$

$\sigma^{ab} = (\gamma^a \gamma^b - \gamma^b \gamma^a)/4$, where $a, b = 0, 1, \dots, d-1$ are the tangent space indices and γ^a are the gamma matrices $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$. The e_μ^a and ω_μ^{ab} are the vierbein and Lorentz connection of anti-de Sitter spacetime. Concrete representation for matrix M may be found in ([12]). As usual it is convenient to introduce generating function for Rarita-Schwinger tensor-spinor field

$$|\psi\rangle = M^{-1} \prod_{l=1}^\nu P_l^{m'_l} |\Psi\rangle, \quad P_l \equiv a_l^b e_b^\mu y_\mu^A \bar{a}_l^A, \quad (43)$$

where a_l^b are new creation operator $[\bar{a}_l^a, a_j^b] = \delta_{ij} \eta^{ab}$, $\eta^{ab} = (-, +, \dots, +)$. Note that the $|\psi\rangle$ is actually a generating function for tangent space Rarita-Schwinger field: $\psi_{a\dots} = e_a^\mu \dots \psi_{\mu\dots}$. Now making use of (42) the equations of motion (30) can be rewritten in terms of $|\psi\rangle$ as follows

$$(\gamma^a e_a^\mu D_{\mu L} + E_0 + \frac{1-d}{2}) |\psi\rangle = 0, \quad (44)$$

$$D_{\mu L} \equiv \partial_\mu + \frac{1}{2} \omega_\mu^{ab} M_{ab}, \quad M^{ab} \equiv \sum_{l=1}^\nu (a_l^a \bar{a}_l^b - a_l^b \bar{a}_l^a) + \sigma^{ab}.$$

In order to transform the gauge transformation to intrinsic coordinates we introduce the gauge parameter $|\lambda_k\rangle$ by analogy with (43)

$$|\lambda_k\rangle = M^{-1} \prod_{l=1}^{\nu} P_l^{m'_{l(k)}} |\Lambda_k\rangle, \quad (45)$$

and now the gauge transformation (33) is rewritten as

$$\delta_{(k)}|\psi\rangle = \sum_{j=0}^{k-1} (-)^j \sum_{l_1 \dots l_{j+1}=1}^{\nu} \delta_{kl_{j+1}} \prod_{i=1}^j \frac{\varepsilon^{l_i l_{i+1}}(a_{l_{i+1}} \bar{a}_{l_i})}{k+1-l_i} a_{l_1}^b (e_b^\mu D_{\mu L} + \frac{1}{2} \gamma_b) |\lambda_k\rangle, \quad (46)$$

where $(a_i a_j) \equiv \eta_{ab} a^a a^b$. Note that the $|\lambda_n\rangle$ satisfies the equation of motion which is obtainable from (44) by making there the substitutions $|\psi\rangle \rightarrow |\lambda_k\rangle$ and $E_0 \rightarrow E_0^\Lambda$. The constraints (6), (8) take the form

$$\bar{a}_n^b (e_b^\mu D_{\mu L} + \frac{1}{2} \gamma_b) |\psi\rangle = 0, \quad \bar{a}_n^b \gamma_b |\psi\rangle = 0, \quad (47)$$

$n = 1, \dots, \nu$, and similar constraints for $|\lambda_k\rangle$ are obtainable from (47) by making there the substitution $|\psi\rangle \rightarrow |\lambda_k\rangle$. The constraints (10) transform to

$$((a_i \bar{a}_i) - m'_i) |\psi\rangle = 0, \quad (\bar{a}_i \bar{a}_i) |\psi\rangle = 0, \quad \varepsilon^{ij} (a_i \bar{a}_j) |\psi\rangle = 0, \quad (48)$$

$i, j = 1, \dots, \nu$, while similar constraints for $|\lambda_k\rangle$ are obtainable from (48) by making there the substitutions $|\psi\rangle \rightarrow |\lambda_k\rangle$ and $m'_i \rightarrow m'_{i(k)}$.

In conclusion let us summarize the results of this letter. For massless fermionic field $|\Psi\rangle$ of arbitrary spin labeled by $\mathbf{m}$ we have constructed: (i) free wave equations of motion in $SO(d-1, 2)$ covariant form (30) as well as in terms of intrinsic coordinates (44); (ii) corresponding gauge transformations (33), (46), subsidiary conditions (6)-(8), (10), (16), (47), (48) and equations of motion for gauge parameter (35) field written also in both forms; (iii) the new representation for equations of motion (39), (40) and gauge transformations (see (33), (41)) in terms of the generators of the anti-de Sitter group $SO(d-1, 2)$; (iv) lowest energy values for massless fermionic field (28) and for gauge parameter field (36). Also we have demonstrated that the gauge parameter for massless field associated with level $k = 1$ Young tableaux is a massive field while for $k > 1$ the gauge parameter is a massless field.

The author would like to thank Prof. B. de Wit for hospitality at Institute for Theoretical Physics of Utrecht University where a part of this work was carried out. This work was supported in part by INTAS contracts No. CT93-0023 and No.96-538, by the Russian Foundation for Basic Research, Grant No.96-01-01144, and by the NATO Linkage, Grant No.931717.

References

- [1] C. Fronsdal, *Rev. Mod. Phys.* **37** (1965) 221; *Phys. Rev.* **D12** (1975) 3819; *Phys. Rev.* **D20** (1979) 848
- [2] S.J. Avis, C.J. Isham and D. Storey, *Phys. Rev.* **D10** (1978) 3565
- [3] K.S. Stelle and P.C. West, *Phys. Rev.* **D21** (1980) 1466
- [4] P.K. Townsend and P. van Nieuwenhuizen, *Phys. Lett.* **B125** (1983) 41
- [5] D.Z. Freedman and G.W. Gibbons, *Nucl. Phys.* **B233** (1984) 24
- [6] E.S. Fradkin and M.A. Vasiliev, *Ann. Phys.* **177** (1987) 63
- [7] M.A. Vasiliev, Higher-spin gauge theories in four, three and two dimensions, Preprint FIAN/TD/24-96, hep-th/9611024, Invited talk at the Quantum Gravity Seminar in memory of M.Markov, June 12-19, 1995 Moscow
- [8] H. Nicolai, Representations of supersymmetry in anti-de Sitter space, in: *Supersymmetry and supergravity*, '84, Ed. B. de Wit, P. Fayet and P. van Nieuwenhuizen (World Scientific, Singapore, 1984)
- [9] M.A. Vasiliev, *Phys. Lett.* **B243** (1990) 378; *Class. Quantum Grav.* **8** (1991) 1387
- [10] M.A. Vasiliev, *Phys. Lett.* **B257** (1991) 111
- [11] C. Fronsdal and R.B. Haugen, *Phys. Rev.* **D12** (1975) 3810
- [12] J. Fang and C. Fronsdal, *Phys. Rev.* **D22** (1980) 1361
- [13] P. Breitenlohner and D.Z. Freedman, *Ann. Phys.* **144** (1982) 249
- [14] M.A. Vasiliev, *Nucl. Phys.* **B301** (1988) 26
- [15] R.R. Metsaev, *Class. Quantum Grav.* **11** (1994) L141
- [16] R.R. Metsaev, *Class. Quantum Grav.* **14** (1997) L115
- [17] R.R. Metsaev, *Phys. Lett.* **B354** (1995) 78
- [18] A.O. Barut and R. Raczka, *Theory of Group Representations and Applications* (PWN-Polish Scientific Publishers, Warszawa 1977)
- [19] D. Zhelobenko, *Uspekhi Mat. Nauk* **17** (1962) 27
- [20] P.A.M. Dirac, *Ann. Math.* **36** (1935) 657
- [21] R.R. Metsaev, *Mod. Phys. Lett.* **A10** (1995) 1719
- [22] S. Deser, J. Kay and K. Stelle, *Phys. Rev.* **D16** (1977) 2445